

TWIST-3 EFFECTS IN POLARIZED PHOTON STRUCTURE *

HIDESHI BABA[†]

*Graduate School of Human and Environmental Studies, Kyoto University
 Kyoto, 606-8501, Japan*

KEN SASAKI[‡]

*Dept. of Physics, Faculty of Engineering, Yokohama National University
 Yokohama, 240-8501, Japan*

TSUNEO UEMATSU[§]

*Dept. of Fundamental Sciences, FIHS, Kyoto University
 Kyoto, 606-8501, Japan*

The polarized photon structure is described by two spin structure functions g_1^γ and g_2^γ which can be studied in the future polarized ep or e^+e^- colliders. Here we investigate the QCD twist-3 effects in g_2^γ to the leading order in QCD.

1. Introduction

In recent years there has been growing interest in the polarized photon structure functions. Especially the first moment of a photon structure function g_1^γ has attracted much attention in connection with the axial anomaly which is also relevant in the nucleon spin structure function $g_1^{p(n)}$. The polarized structure function g_1^γ could be experimentally studied in the polarized version of ep collider HERA,^{1,2} or more directly measured by the polarized e^+e^- collision in the future linear collider. And the next-to-leading order QCD analysis of g_1^γ has been recently performed in the literature.^{3,4,5}

Now there exists another structure function g_2^γ for the virtual photon target, where the twist-3 effect is also relevant in addition to the usual twist-2 effect. In this talk, we investigate the twist-3 effects in g_2^γ , to the leading order in QCD, and show that it is a sizable effect in contrast to the nucleon case,⁶ where it appears to be small in the experimental data.^{7,8}

*Presented by T. Uematsu at the 3rd Circum-Pan-Pacific Symposium on “High Energy Spin Physics”, Beijing, October 8-13, 2001. KUCP-204, YNU-HEPTh-02-104, to appear in the Proceedings.

[†]e-mail address: baba@phys.h.kyoto-u.ac.jp

[‡]e-mail address: sasaki@cnb.phys.ynu.ac.jp

[§]e-mail address: uematsu@phys.h.kyoto-u.ac.jp

2. Photon Structure Function $g_2^\gamma(x, Q^2, P^2)$ and Twist-3 Effects

Let us consider the polarized deep inelastic scattering on a polarized virtual photon target and study the virtual photon structure functions for the kinematical region:

$$\Lambda^2 \ll P^2 \ll Q^2, \quad (1)$$

where $q^2 = -Q^2$ ($p^2 = -P^2$) is the mass squared of the probe (target) photon, and Λ is the QCD scale parameter.

The antisymmetric part of the structure tensor, $W_{\mu\nu\rho\tau}^A(p, q)$ for the target (probe) photon with momentum p (q), relevant for the polarized structure, can be written in terms of the structure functions, g_1^γ and g_2^γ , as ⁴

$$W_{\mu\nu\rho\tau}^A = \frac{1}{(p \cdot q)^2} \left[p \cdot q \epsilon_{\mu\nu\lambda\sigma} \epsilon_{\rho\tau}^{\sigma\beta} q^\lambda p_\beta g_1^\gamma - (\epsilon_{\mu\nu\lambda\sigma} \epsilon_{\rho\tau\alpha\beta} q^\lambda p^\sigma q^\alpha p^\beta - p \cdot q \epsilon_{\mu\nu\lambda\sigma} \epsilon_{\rho\tau}^{\sigma\beta} q^\lambda p_\beta) g_2^\gamma \right] \quad (2)$$

Here we note that we have the structure function g_2^γ only for non-zero P^2 , i.e. virtual photon.

Now we decompose g_2 into twist-2 and twist-3 contributions:

$$g_2 = g_2^{\text{tw.2}} + g_2^{\text{tw.3}}. \quad (3)$$

where $g_2^{\text{tw.2}}$ can be written by using Wandzura-Wilczek relation:⁹

$$g_2^{\text{tw.2}} = g_2^{\text{WW}}(x, Q^2) \equiv -g_1(x, Q^2) + \int_x^1 \frac{dy}{y} g_1(y, Q^2). \quad (4)$$

Experimental data for the nucleon show $g_2 \approx g_2^{\text{tw.2}} = g_2^{\text{WW}}$, and now we ask what about the photon structure, especially virtual photon case ?

3. OPE and QCD Effects

First we consider the operator product expansion (OPE) relevant for the photon structure functions. The OPE sandwiched between the photon states can be decomposed into the twist-2 and twist-3 contributions as follows:

$$\begin{aligned} & \int d^4x e^{iq \cdot x} \langle \gamma(p, s) | J_\mu(x) J_\nu(0)^{[A]} | \gamma(p, s) \rangle \\ & \sim \sum_n E_n^{(2)}(Q^2) \langle \gamma(p, s) | R_n^{(2)} | \gamma(p, s) \rangle + \sum_n E_n^{(3)}(Q^2) \langle \gamma(p, s) | R_n^{(3)} | \gamma(p, s) \rangle. \end{aligned} \quad (5)$$

where $R_n^{(2)}$ and $R_n^{(3)}$ denote the twist-2 and twist-3 operators, respectively. For the nucleon the matrix element: $\langle N(p, s) | R_n^{(3)} | N(p, s) \rangle$ is small. Now we calculate the virtual photon structure functions arising from the so-called Box diagram, with $\langle e^4 \rangle = \sum_{i=1}^{N_f} e_i^4 / N_f$, and N_f being the number of active flavors. It turns out that the twist-3 contribution to g_2^γ is actually non-vanishing: We have plotted the g_1^γ , g_2^γ and $\bar{g}_2^\gamma$ as a function of x for the virtual photon target, where $Q^2 = 30 \text{ GeV}^2$ and $P^2 = 1 \text{ GeV}^2$ for $N_f = 3$ in Fig.1.




Fig. 1. The Box-diagram contributions to $g_1^\gamma(x, Q^2, P^2)$ (dashed), $g_2^\gamma(x, Q^2, P^2)$ (solid) and $\bar{g}_2^\gamma(x, Q^2, P^2)$ (dash-2dotted) for $Q^2 = 30 \text{ GeV}^2$ and $P^2 = 1 \text{ GeV}^2$ for $N_f = 3$.

Now the n -th moment of the twist-3 part $\bar{g}_2^\gamma \equiv g_2^\gamma - g_2^{\gamma(WW)}$ reads

$$\int_0^1 dx x^{n-1} \bar{g}_2^\gamma(x, Q^2, P^2) \approx \frac{n-1}{2n} \vec{X}_n(Q^2/P^2, \bar{g}(Q^2), \alpha) \vec{E}_n(1, \bar{g}(Q^2)) \quad (6)$$

in the leading logarithmic order (LO) in QCD, where

$$\vec{X}_n = \int_{\bar{g}}^g dg \frac{\vec{K}_n(g, \alpha)}{\beta(g)} T \exp \left[\int_{\bar{g}}^g dg' \frac{\hat{\gamma}_n(g')}{\beta(g')} \right] \quad (7)$$

with $\hat{\gamma}_n$ representing the mixing of the twist-3 hadronic operators and $\vec{K}_n$ describing the mixing between twist-3 hadronic and photonic operators. Thus, the flavor non-singlet (NS) part reads

$$\begin{aligned} \int_0^1 dx x^{n-1} \bar{g}_2^{\gamma(NS)}(x, Q^2, P^2) &= \frac{n-1}{n} \frac{\alpha}{4\pi} \cdot \frac{1}{2\beta_0} (-24N_f) (\langle e^4 \rangle - \langle e^2 \rangle^2) \frac{1}{n(n+1)} \\ &\times \frac{1}{1 + \lambda_{NS}^n/2\beta_0} \frac{4\pi}{\alpha_s(Q^2)} \left\{ 1 - \left(\frac{\alpha_s(Q^2)}{\alpha_s(P^2)} \right)^{\lambda_{NS}^n/2\beta_0+1} \right\} \end{aligned}$$

In the large N_c limit ($N_c \rightarrow \infty$) we have the 1-loop anomalous dimension of the NS twist-3 operator:^{10,11}

$$\lambda_{NS}^n = \gamma_{(3)NS}^{0,n} = 8C_F \left(S_n - \frac{1}{4} - \frac{1}{2n} \right) \quad \text{where} \quad S_n = \sum_{j=1}^n \frac{1}{j} \quad (8)$$

up to $1/N_c^2$. We have plotted in Fig.2 the above result to LO in QCD for the flavor non-singlet part. We expect the similar behavior for the flavor-singlet contribution.




Fig. 2. The Box-diagram (dashed) and the QCD LO (solid) contributions to $\bar{g}_2^\gamma(x, Q^2, P^2)$ for $Q^2 = 30 \text{ GeV}^2$ and $P^2 = 1 \text{ GeV}^2$ for $N_f = 3$.

4. Concluding Remarks

We have investigated the twist-3 effect in g_2^γ for the virtual photon target, in the LO QCD and have found it gives rise to a sizable effect. More thorough QCD analysis including flavor-singlet part is now under investigation.

Acknowledgements

One of the authors (T.U.) thanks the organizers of the symposium, especially to Prof. Ma for the wonderful organization and the hospitality during the symposium.

References

1. D. Barber, in proceedings of the “Zeuthen Workshop on the Prospects of Spin Physics at HERA”, DESY 95-200, p.76, eds. J. Blümlein and W. D. Nowak.
2. M. Stratmann and W. Vogelsang, *Z. Phys.* **C74**, 641 (1997).
3. M. Stratmann and W. Vogelsang, *Phys. Lett.* **B386**, 370 (1996).
4. K. Sasaki and T. Uematsu, *Phys. Rev.* **D59**, 114011 (1999); *Phys. Lett.* **B473**, 309 (2000); *Eur. Phys. J.* **C20**, 283 (2001).
5. M. Glück, E. Reya and C. Sieg, *Phys. Lett.* **B503**, 285 (2001); *Eur. Phys. J.* **C20**, 271 (2001).
6. R. L. Jaffe, these proceedings.
7. E143 Collaboration, K. Abe et al., *Phys. Rev. Lett.* **76**, 587 (1996).
8. E155 Collaboration, P. L. Anthony et al., *Phys. Lett.* **B458**, 529 (1999).
9. S. Wandzura and F. Wilczek, *Phys. Lett. B* **172**, 195 (1977).
10. A. Ali, V. M. Braun and G. Hiller, *Phys. Lett.* **B266**, 117 (1991).
11. K. Sasaki, *Phys. Rev.* **D58**, 094007 (1998).